TRIP6 Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID NP_003293.1

Catalog No. A300-865A GeneID 7205

Lot No. A300-865A-1

APPLICATIONS WB, IP
SPECIES REACTIVITY Human

PRESUMED REACTIVITY Based on 100% sequence identity, this antibody is predicted to react with Bovine

AMOUNT 100 μl

CONCENTRATION 200 μg/ml

STORAGE/SHELF LIFE 2 – 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-buffered Saline containing 0.1% BSA and 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION PROCEDURES

Antibody was affinity purified using an epitope specific to TRIP6 immobilized on solid support.

The epitope recognized by A300-865A maps to a region between residue 1 and 50 of human Thyroid Hormone Receptor Interactor 6 using the numbering given in entry NP_003293.1

(GeneID 7205).

Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280 nm

of 1.4 equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:2,000 - 1:10,000

Immunoprecipitation 2 – 5 µg/mg lysate

APPLICATION NOTES Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020),

Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-20% SDS-PAGE

(link to IP-western blot protocol in Additional Info section below).

Western blot of lysates performed using standard western blot reagents and 4-20% SDS-PAGE.

ADDITIONAL INFO https://www.bethyl.com/product/A300-865A

Use the link above to view SDS, a current list of citations, and other product specific information.


IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer

Date: June 21, 2019



TRIP6 Antibody A300-865A



Detection of human TRIP6 by western blot and immunoprecipitation. Samples: Whole cell lysate from HeLa (5, 15 and 50 μg for WB; 1 mg for IP, 20% of IP loaded) and HEK293T (T; 50 μg) cells. Antibodies: Affinity purified rabbit anti–TRIP6 antibody BL1858 (Cat. No. A300–865A) used for WB at 0.04 μg/ml (A) and 1 μg/ml (B) and used for IP at 3 μg/mg lysate (B). TRIP6 was also immunoprecipitated by rabbit anti–TRIP6 antibodies BL1859, BL452 (Cat. No. A300–334A) and BL453 (Cat. No. A300–335A), which recognize other epitopes. Detection: Chemiluminescence with exposure times of 3 minutes (A) and 10 seconds (B).