YKT6 Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID 015498.1 Catalog No. A305-479A GeneID 10652

Lot No. A305-479A-1

APPLICATIONS WB, IP

SPECIES REACTIVITY Human, Mouse

PRESUMED REACTIVITY Based on 100% sequence identity, this antibody is predicted to react with Bovine

AMOUNT 100 μl

CONCENTRATION 1000 μg/ml

STORAGE/SHELF LIFE 2 – 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION PROCEDURES

Antibody was affinity purified using an epitope specific to YKT6 immobilized on solid support.

The epitope recognized by A305–479A maps to a region between residue 125 to 175 of human Synaptobrevin homolog YKT6 using the numbering given in entry O15498.1 (GenelD 10652).

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4

equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:500 - 1:2.500

Immunoprecipitation 2 – 10 µg/mg lysate

APPLICATION NOTES Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. \$100-020),

Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-12% SDS-PAGE

(link to IP-western blot protocol in Additional Info section below).

Western blot of lysates performed using standard western blot reagents and 4-12% SDS-PAGE.

ADDITIONAL INFO https://www.bethyl.com/product/A305-479A

Use the link above to view SDS, a current list of citations, and other product specific information.


IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB


This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer

Date: June 21, 2019



YKT6 Antibody A305-479A





Detection of human and mouse YKT6 by western blot. Samples: Whole cell lysate (15 μ g) from HeLa, HEK293T, Jurkat, mouse TCMK-1, and mouse NIH 3T3 cells prepared using NETN lysis buffer. Antibody: Affinity purified rabbit anti-YKT6 antibody A305-479A (lot A305-479A-1) used for WB at 0.1 μ g/ml. Detection: Chemiluminescence with an exposure time of 3 minutes.

Detection of human YKT6 by western blot of immunoprecipitates. Samples: Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) from 293T cells prepared using NETN lysis buffer. Antibodies: Affinity purified rabbit anti-YKT6 antibody A305-479A (lot A305-479A-1) used for IP at 6 μg per reaction. YKT6 was also immunoprecipitated by rabbit anti-YKT6 antibody A305-480A. For blotting immunoprecipitated YKT6, A305-479A was used at 1 μg/ml. Detection: Chemiluminescence with an exposure time of 30 seconds.