

 Phone:
 888-558-5227

 651-644-8424

 Fax:
 888-558-7329

 Email:
 getinfo@lktlabs.com

 Web:
 lktlabs.com

Product Information

Product ID	D1759			
CAS No.	51481-10-8		H O H	
Chemical Name				
Synonym	Vomitoxin	O Ū Ū		
Formula	C ₁₅ H ₂₀ O ₆			
Formula Wt.	296.32			
Melting Point	151-153°C	Bulk quanitites available upon request		
Purity	≥98%, TLC	Product ID	Size	
Solubility	Soluble in acetonitrile, methanol and ethyl	D1759 D1759	1 mg 5 mg	
Store Temp	4°C			
Ship Temp	Ambient			

escription Deoxynivalenol is a trichothecene mycotoxin initially produced by species of *Fusarium* that is found in cereals and other products of wheat or similar grains. Deoxynivalenol exhibits immunomodulatory, cytotoxic, and pro-inflammatory activities. In vivo, deoxynivalenol increases the formation of pores in the intestinal epithelial barrier of the jejunum, increases the number of CD16+ cells, and downregulates expression of syndeca, fibulin 6, and BM-40. In vitro, this toxin activates p38 MAPK and p53 to induce activation of caspases 3, 8, and 9, resulting in apoptosis; it also induces rRNA cleavage, inhibiting ribosomal translation. In other cellular models, deoxynivalenol upregulates expression of IL-18, IL-6, TNF-α, COX-2, and mPGES-1.

References Ovando-Martínez M, Ozsisli B, Anderson J, et al. Analysis of deoxynivalenol and deoxynivalenol-3-glucoside in hard red spring wheat inoculated with Fusarium graminearum. Toxins (Basel). 2013 Dec 17;5(12):2522-32. PMID: 24351715.

Nossol C, Diesing AK, Kahlert S, et al. Deoxynivalenol affects the composition of the basement membrane proteins and influences en route the migration of CD16(+) cells into the intestinal epithelium. Mycotoxin Res. 2013 Nov;29(4):245-54. PMID: 23949948.

He K, Zhou HR, Pestka JJ. Targets and intracellular signaling mechanisms for deoxynivalenol-induced ribosomal RNA cleavage. Toxicol Sci. 2012 Jun;127(2):382-90. PMID: 22491426.

Girardet C, Bonnet MS, Jdir R, et al. Central inflammation and sickness-like behavior induced by the food contaminant deoxynivalenol: a PGE2-independent mechanism. Toxicol Sci. 2011 Nov;124(1):179-91. PMID: 21873375.

Caution: This product is intended for laboratory and research use only. It is not for human or drug use.