



LKT Laboratories, Inc.
Hexamethylene Bisacetamide

Phone: 888-558-5227
651-644-8424
Fax: 888-558-7329
Email: getinfo@lktlabs.com
Web: lktlabs.com

Product Information

Product ID H1892

CAS No. 3073-59-4

Chemical Name N,N'-Hexamethylene (bis)acetamide

Synonym N,N'-Diacetyl-1,6-hexanediamine, HMBA


Formula C₁₀H₂₀N₂O₂

Formula Wt. 200.28

Melting Point 128-129°C

Purity ≥98%

Solubility Soluble in water or ethanol.



Bulk quantities available upon request

Product ID	Size
H1892	25 g
H1892	50 g

Store Temp Ambient

Ship Temp Ambient

Description Hexamethylene bisacetamide (HMBA) activates hexamethylene bisacetamide-inducible protein 1 (HEXIM1), inhibiting transcription elongation factor b (P-TEFb), a regulator of RNA polymerase II and mRNA synthesis. HMBA exhibits anticancer activity and is used to induce cellular differentiation and apoptosis in cancer cells. HMBA also inhibits activation of NF-κB, Akt, and ERK.

References Lew QJ, Chia YL, Chu KL, et al. Identification of HEXIM1 as a positive regulator of p53. J Biol Chem. 2012 Oct 19;287(43):36443-54. PMID: 22948151.

Dey A, Wong E, Kua N, et al. Hexamethylene bisacetamide (HMBA) simultaneously targets AKT and MAPK pathway and represses NF kappaB activity: implications for cancer therapy. Cell Cycle. 2008 Dec;7(23):3759-67. PMID: 19029824.

Zhang Z, Liong EC, Lau TY, et al. Induction of apoptosis by hexamethylene bisacetamide is p53-dependent associated with telomerase activity but not with terminal differentiation. Int J Oncol. 2000 May;16(5):887-92. PMID: 10762623.

Caution: This product is intended for laboratory and research use only. It is not for human or drug use.